

Table 2. Main kinematics data

ident.	other ident.	V_{hel} km s $^{-1}$	σ_0 km s $^{-1}$	Type
NGC 2300		2003±34	330(1)	E2
HCG 62a		4279±25	246±24	E3
HCG 62b		3542±25	272±26	S0
HCG 68a	NGC 5353	2250±22	361±22	S0
HCG 68b	NGC 5354	2570±17	241±22	E2
K 416b	NGC 5481	2033±13	185±14	E3

Notes: (1) The central velocity dispersion of NGC 2300 is extrapolated.

Table 1. ASCA upper limits of X-ray emission from K 416 members

Object	counts s ⁻¹ (10 ⁻³)	Flux _(0.3–10keV) (10 ⁻¹³ ergs s ⁻¹ cm ⁻²)	L _(0.3–10keV) (10 ⁴¹ h ₅₀ ⁻¹ ergs s ⁻¹)	Flux _(0.3–2.4keV) (10 ⁻¹³ ergs s ⁻¹ cm ⁻²)	L _(0.3–2.4keV) (10 ⁴¹ h ₅₀ ⁻¹ ergs s ⁻¹)	detector
NGC 5480	< 1.4	< 1.0	< 0.2	< 0.5	< 0.1	GIS 2
NGC 5480	< 3.0	< 1.7	< 0.3	< 0.9	< 0.2	GIS 3
NGC 5481	< 1.4	< 0.9	< 0.2	< 0.8	< 0.2	GIS 2
NGC 5481	< 2.6	< 1.4	< 0.3	< 1.2	< 0.3	GIS 3
Extended Emission	< 3.3	< 2.0	< 0.4	< 1.9	< 0.4	GIS 2
	< 5.0	< 2.6	< 0.5	< 2.4	< 0.5	GIS 3

Notes: 3 σ upper limits have been calculated in a circle of 2'5 for the two galaxies and of 7'5 for the extended emission, as explained in the text. Fluxes are derived assuming a thermal bremsstrahlung model with $kT=5$ keV for NGC 5480 and a Raymond Smith plasma model with $kT=1$ keV and abundance = 0.5 solar for NGC 5481 and the extended source. N_H has been taken from Stark et al. (1992) = 1.41×10^{20} cm⁻². A distance of 40 h₅₀⁻¹ Mpc is assumed.

Table 3. Positions and parameters of the fit of X-ray spectra of unknown sources

Object	R.A. (2000)	δ (2000)	Counts/s (10 ⁻³)	γ	norm. (10 ⁻⁵)	χ^2	bins	Flux _(0.3–10keV) (10 ⁻¹³ erg cm ⁻² s ⁻¹)	Camera
Unknown source 1	14 05 12	50 42 29							GIS 2
	14 05 04	50 41 36	4.4±0.5	2.3 ^{+2.0} _{-0.8}	9.4	0.8	15	4.7	GIS 3
Unknown source 2	14 05 37	50 56 51							GIS 2
	14 05 29	50 56 11	4.0±0.4	1.8 ^{+0.9} _{-0.3}	6.8	0.8	16	3.7	GIS 3

Notes: GIS 3 spectra were tentatively fitted with a power law, γ , and interstellar absorption, $N_H = 0$ and $1.1 \cdot 10^{21}$ cm⁻² respectively for the source 1 and 2. The errors are computed on two degrees of freedom 90% confidence interval.